

Track 1A Cardiology Critical Care



Critical Care Cardiology Session American College of Cardiology Kentucky Chapter



Session

Moderator: Jasdeep Dhaliwal, MD

Speakers: Vasili Katsadouros, DO, Sonu Abraham, MD, Siddarth Pahwa, MD, and Jeffrey

Spindel, DO

Session Chair: Paul Anaya, MD PhD



Agenda

2:05 PM Introduction

2:10-2:15 PM Case Presentation: A Shocking Success, Vasili Katsadouros, DO

2:15-2:25 PM Approach to the Patient with Cardiogenic Shock, Sonu Abraham, MD

2:30-2:40 PM Cardiogenic Shock: A Surgeon's Perspective, Siddharth Pahwa, MD

2:45-2:55 PM The Future of Critical Care Cardiology



Trainee Presentation; Vasili Katsadouros




A Shocking Success KY-ACC Critical Care

Vasili Katsadouros DO



Our Case





Over last year, multiple symptoms limiting ADL/iADL Week prior, symptoms of URI



Presented to outside hospital with abdominal pain
Initial presentation concerning for shock

LVEF found to be under 20%



Required multiple inotropes and vasopressors

Refractory shock despite above interventions



She was then transferred to UK for further management

Clinical picture concerning for cardiogenic shock



Our Case


RHC

- RA 18,
- PA 35/20/28,
- PAWP 19,
- PA sat 56%,
- CO/CI Fick 3.95/2.3,
- PVR 2.25, SVR 1280 dynes,
- PAPi 0.83
- CPO 0.73
- CVP to Wedge 0.95

| <u>Cardiogenic Shock Stage</u> | Study Definition |
|------------------------------------|-------------------------------------------------------|
| Stage A (" <u>A</u> t risk") | Neither hypotension/tachycardia nor hypoperfusion |
| Stage B (" <u>B</u> eginning") | Hypotension/tachycardia WITHOUT hypoperfusion |
| Stage C ("Classic") | Hypoperfusion WITHOUT deterioration |
| Stage D (" <u>D</u> eteriorating)" | Hypoperfusion WITH deterioration NOT refractory shock |
| Stage E (" <u>E</u> xtremis") | Hypoperfusion WITH deterioration AND refractory shock |





Our Case





EMB Results – Lymphocytic Myocarditis








Our Case – Unfolding the story

- Lymphocytic myocarditis
- Respiratory panel (+) Rhinovirus
- Questionable underlying chronic cardiac dysfunction






Our Case – Multi-disciplinary care

- Inability to wean support
- Permanent MCS VS Transplant
 - Potential for myocardial recovery
 - Young age
 - Positive QuantiFeron TB gold test
- Decision made for LVAD implantation

Prospective Multicenter Study of Myocardial Recovery Using Left Ventricular Assist Devices (RESTAGE-HF [Remission from Stage D Heart Failure])


Medium-Term and Primary End Point Results






Our Case — Post VAD

- Post op course
 - -Complicated by...
 - Significant bleeding
 - RV failure
- Genetic Testing
 - HCM/DCM variant
- Myocardial recovery regimen
- Discharged home and doing well!



HeartMate 3 LVAD



- Intra-thoracic pump
- Improved coagulation system stability
- Fully magnetically levitated motor
- Artificial pulsatility
- Large blood flow pathways



Successful management of cardiogenic shock



References

Senman B, Jentzer JC, Barnett CF, et al. Need for a Cardiogenic Shock Team Collaborative-Promoting a Team-Based Model of Care to Improve Outcomes and Identify Best Practices. *J Am Heart Assoc.* 2024;13(6):e031979. doi:10.1161/JAHA.123.031979

Jentzer JC, van Diepen S, Barsness GW, et al. Cardiogenic Shock Classification to Predict Mortality in the Cardiac Intensive Care Unit. *J Am Coll Cardiol*. 2019;74(17):2117-2128. doi:10.1016/j.jacc.2019.07.077

Varshney AS, DeFilippis EM, Cowger JA, Netuka I, Pinney SP, Givertz MM. Trends and Outcomes of Left Ventricular Assist Device Therapy: JACC Focus Seminar. *J Am Coll Cardiol*. 2022;79(11):1092-1107. doi:10.1016/j.jacc.2022.01.017



Talk 1: Sonu Abraham



Cardiogenic Shock: An Update

Sonu Abraham, MD, FACC
Advanced Heart Failure & Transplant Cardiologist
Assistant Professor, Department of Medicine
Gill Heart & Vascular Institute
University of Kentucky



Learning Objectives

Cardiogenic shock Definition, Staging and Classification

Pharmacological therapies for cardiogenic shock

Invasive therapies for cardiogenic shock


Utility of the Swan-Ganz Catheter



Cardiogenic Shock Definition, Staging and Classification




Cardiogenic Shock Spiral





van Diepen S et al. *Circulation*. 2017;136(16):e232-e268.

Heterogeneity of Cardiogenic Shock






Hemodynamic presentations of Cardiogenic Shock

| | | Volume Status | | |
|-------------|------|---------------------------------------------------------------------|-----------------------------|--|
| 27 | | Wet | Dry | |
| Ĕ | | Classic Cardiogenic Shock | Euvolemic Cardiogenic Shock | |
| Circulation | Cold | (↓CI; 个SVRI; 个PCWP) | (↓CI; ↑SVRI; ↔PCWP) | |
| | | Vasodilatory Cardiogenic Shock | Vasodilatory Shock | |
| era | | or | (Not Cardiogenic Shock) | |
| Periphera | Warm | Mixed Shock | | |
| | | $(\downarrow CI; \downarrow / \leftrightarrow SVRI; \uparrow PCWP)$ | (个CI; ↓SVRI; ↓PCWP) | |




van Diepen S et al. Circulation. 2017;136(16):e232-e268.

Modified SCAI Shock Stages





Short-term mortality as a function of SCAI SHOCK stages






Naidu, Srihari S. et al. JSCAI, Volume 1, Issue 1, 100008

Cardiac Arrest as a Shock Modifier

Jentzer, J et al. JACC. 2019 Oct, 74 (17) 2117–2128





3-axis Model of Cardiogenic Shock Evaluation & Prognostication

Naidu, Srihari S. et al. JSCAI, Volume 1, Issue 1, 100008



Shock severity SCAI shock stage Hemodynamics Metabolic derangements Vasopressor toxicity Phenotype & Etiology Acute vs. Acute-on-Cardiogenic Chronic shock patient Clinical etiology of CS **Risk modifiers** RV vs. LV vs. BIV systolic Non-modifiable risk dysfunction factors: e.g. age, Cardiac vs. comorbidities Risk modifiers cardiopulmonary failure Cardiac arrest with coma Congestion profile Presence/reversibility Biochemical phenotype of organ failure Systemic inflammatory response Frailty/risk of complication

Chronic cardiovascular disease SCAI shock stage not applicable Recovery RECOVERY PATHWAYS

Normalization of perfusion metrics while on support (MCS or pharmacologic) improves to Stage C. If remains normal with removal of support, then improves to Stage B or A.



Hemodynamically <u>stable</u> SCAI shock stage A

Loss of ‡ compensation

Hemodynamically <u>unstable</u>

SCAI shock stage B

Loss of **‡** compensation

Hypoperfusion = Shock SCAI shock stage C

Deterioration

Failure to stabilize with initial Tx SCAI shock stage D

1 Deterioration

Extremis / refractory shock
SCAI shock stage E


DETERIORATION PATHWAYS


Acute catastrophic event (i.e., prolonged CA) arrives in Stage E. All others must stop at least transiently in Stage C for first intervention.



Naidu, Srihari S. et al. JSCAI, Volume 1, Issue 1, 100008

Dynamic Escalation of CS stages








SCAI-CSWG Shock Stages


Kapur NK et al. *J Am Coll Cardiol*. 2022;80(3):185-198



CENTRAL ILLUSTRATION: Clinical Variables and Parameters to Define Society for Cardiovascular Angiography and Interventions Stages



Shock Academic Research Consortium (SHARC) Classification of Cardiogenic Shock





Waksman R et al. *Circulation*. 2023;148(14):1113-1126.

SHARC Classification of Cardiogenic Shock

AMI-CS

 In setting of acute MI (STEMI or NSTEMI) in which acute ischemia is the primary driver of cardiogenic shock.

STEMI

NSTEMI

HF-CS

 Primary ventricular (L/R) failure (in absence of acute MI) De novo HF-CS

Acute-on-chronic

Sub-classify by

specific cause

Specific Etiologies

Key Specific etiologies:

Post-cardiac

arrest or eCPR

Additional

LV/RV/BiV

Modifiers

Mycarditis

Takotsubo

Amyloid

Peripartum

 Mixed shock syndrome

Secondary CS

• Due to another primary non-myocardial cardiac cause: arrhythmia, pericardial or valvular disease.

In setting of cardiac surgery


Sub-classify by surgery type

Post-cardiotomy CS

| Kentucky CHAPTER

Waksman R et al. Circulation. 2023;148(14):1113-1126.


Hospital
Mortality as
a Function
of SCAI
Shock Stage



Jentzer, J et al. JACC. 2019 Oct, 74 (17) 2117-2128



De Novo HF-CS vs Acute on Chronic HF-CS





Bhatt AS et al. *J Card Fail*. 2021;27(10):1073-1081

Interim Summary

Of cardiogenic shock etiologies, **HF-CS** now exceeds AMI-CS.

Staging of shock at frequent intervals is important as cardiogenic shock is dynamic (B for Bad!!!)

Congestion is associated with worse outcomes.

Cardiac arrest worsens outcomes in cardiogenic shock.

In-hospital mortality is more in patients with AMI-CS when compared to HF-CS.

Within HF-CS, De-novo HF-CS has higher mortality than Acute on Chronic HF-CS.




Pharmacological Therapy for Cardiogenic Shock



Vasopressors

Which vasopressor do you prefer to use in cardiogenic shock?





Vasopressors


• The initial goal of therapy in patients with CS and associated hypotension should be focused on the restoration of perfusion to vital organs, achieved through augmenting MAP and cardiac output.






Comparison of Dopamine and Norepinephrine in the Treatment of Shock

Authors: Daniel De Backer, M.D., Ph.D., Patrick Biston, M.D., Jacques Devriendt, M.D., Christian Madl, M.D., Didier Chochrad, M.D., Cesar Aldecoa, M.D., Alexandre Brasseur, M.D., Pierre Defrance, M.D., Philippe Gottignies, M.D., and Jean-Louis Vincent, M.D., Ph.D., for the SOAP II Investigators* Author Info & Affiliations








Epinephrine Versus Norepinephrine for Cardiogenic Shock After Acute Myocardial Infarction N= 57



N= 57 patients

Bruno Levy, MD, PhD,^a Raphael Clere-Jehl, MD,^b Annick Legras, MD,^c Tristan Morichau-Beauchant, MD,^d Marc Leone, MD, PhD,^e Ganster Frederique, MD,^f Jean-Pierre Quenot, MD, PhD,^g Antoine Kimmoun, MD, PhD,^a Alain Cariou, MD, PhD,^d Johan Lassus, MD, PhD,^h Veli-Pekka Harjola, MD, PhD,^h Ferhat Meziani, MD, PhD,^b Guillaume Louis, MD,^f Patrick Rossignol, MD, PhD,^f Kevin Duarte, PhD,^f Nicolas Girerd, MD, PhD,^f Alexandre Mebazaa, MD, PhD,^k Philippe Vignon, MD, PhD^l



- (1) There were no differences observed in MAP, cardiac index, or stroke volume in patients treated with either epinephrine or norepinephrine
- (2) Epinephrine treatment was associated with worse metabolic acidosis and increased lactate levels
- (3) Those treated with Epi experienced significantly greater increases in heart rate and a concomitant increase in the cardiac double product, an indirect surrogate of myocardial oxygen consumption
- (4) The study was terminated prematurely due to increased rates of refractory CS (odds ratio [OR], 8.24 [95% CI, 1.61–42.18], P=0.01) developing in the epinephrine-treated arm.



Cochrane analysis

- 19 eligible studies including 2385 individuals
- Insufficient evidence to support the superiority of a particular inotrope or vasodilating agent in cardiogenic shock, especially with respect to a mortality benefit.

| | Epinep | | Norepinephrine-d | | Risk Ratio | Risk Ratio |
|-------------------|--------|-------|------------------|-------|---------------------|-------------------------------|
| Study or Subgroup | Events | Total | Events | Total | M-H, Random, 95% CI | M-H, Random, 95% CI |
| Levy 2011 | 5 | 15 | 4 | 15 | 1.25 [0.41 , 3.77 | 7] |
| | | | | | | 0.01 0.1 1 10 |
| | | | | | Fa | vours Epinephrine Favours Nor |



Uhlig K et al. Cochrane Database of Systematic Reviews 2020, Issue 11. Art. No.: CD009669.

Current guidelines of vasopressor therapy

Indication to commence therapy

Hypotension (systolic blood pressure <90 mm Hg) and/or evidence of end-organ hypoperfusion

Societal guideline

ESC HF

ACC/AHA HF

ESC STEMI

AHA CS

AHA AMI-CS

Recommendation and level of evidence

IIb - B/C

IIa - B

IIb - C

N/A

N/A

First-line agent

Hypotensive: Norepinephrine

Low output: No recommendation

Hypotensive: Clinician preference

Low output: Clinician

preference

Hypotensive: Norepinephrine

Low output: Dobutamine Hypotensive: Norepinephrine or Dopamine

Low output: Norepinephrine or Dopamine Hypotension: Norepinephrine



Bloom JE et al. J Am Heart Assoc. 2023;12(15):e029787.


Inotropes

ORIGINAL ARTICLE



Milrinone as Compared with Dobutamine in the

The NEW ENGLAND JOURNAL of MEDICINE Milrinone vs. Dobutamine in Cardiogenic Shock **DOUBLE-BLIND, RANDOMIZED TRIAL Dobutamine** Milrinone 192 N = 96N = 96Patients admitted to the cardiac ICU with cardiogenic shock 54% In-hospital death from any cause, TIA, stroke, or cardiovascular 52 patients 47 patients or renal events Relative risk, 0.90; 95% CI, 0.69-1.19; P=0.47 No between-group difference was observed in the primary composite outcome or in important secondary outcomes. R. Mathew et al. 10.1056/NEJMoa2026845 Copyright © 2021 Massachusetts Medical Society





Inotropes: Signal towards harm?

Original Contribution



Short-term Intravenous Milrinone for Acute Exacerbation of Chronic Heart Failure

A Randomized Controlled Trial


Michael S. Cuffe, MD; Robert M. Califf, MD; Kirkwood F. Adams, Jr, MD; et al

- Increased in-hospital mortality
- Increased atrial arrhythmias
- Increased hypotension




Early Serial Assessment of Aggregate Vasoactive Support and Mortality in Cardiogenic Shock: Insights From the Critical Care Cardiology Trials Network Registry

Vasoactive-inotropic score (VIS) = dobutamine + dopamine + 10 × phenylephrine + 10 × milrinone + 100 × epinephrine + 100 × norepinephrine + 10 000 × vasopressin




Circulation: Heart Failure • Volume 17, Number 5 • https://doi.org/10.1161/CIRCHEARTFAILURE.124.011736





CAPITAL DOREMI II TRIAL



- Participant admitted to an ICU and diagnosed with SCAI class C or D cardiogenic shock
- Research team member confirms that subject meets all eligibility criteria
- Participant or substitute decision maker reviews and signs consent form
- Treating physician selects their inotrope of choice

Randomization in 1:1 ratio using a centralized, web-based software algorithm

Inotrope Group:

Dobutamine or milrinone as selected by the treating physician

Placebo Group:

0.9% NaCl at an infusion rate comparable to that of the inotrope arm

Intervention Period

- Continued for 12 hours or until participant reaches primary endpoint
- Frequent clinical reassessments with drugs titrated from stages 0 to 5

| Stage | Dobutamine (mcg/kg/min) | Milrinone (mcg/kg/min) | 0.9% NaCl (mL/hour) |
|-------|----------------------------|---------------------------|------------------------|
| 0 | 0 | 0 | 0 |
| 1 | 2.5 | 0.125 | 5 |
| 2 | 5.0 | 0.250 | 10 |
| 3 | 7.5 | 0.375 | 15 |
| 4 | 10 | 0.500 | 20 |
| 5 | >10 | >0.500 | 25 |

Participant Follow-Up

- Following intervention period, participant continues open-label therapy at the discretion of treating physician
- All secondary outcomes assessed at time of discharge from hospital




Congestion matters!


Dynamic Invasive Hemodynamic Congestion Profile Impacts Acute Myocardial Infarction Complicated by Cardiogenic Shock Outcomes: A Real-World Single-Center Study 80 Survival probability (%) Δ Congestion Congestive Decongestive · · · Neutral Log-rank p=0.005 20 Time (days) Worse hemodynamics in SCAI D/E and Biventricular congestion Higher: · Right atrial pressure Cardiac output, power and index PCWP Arterial pressures · Pulmonary pressure 295 patients with AMI-CS, D · Heart rate with early (first 24 h of CS onset) hemodynamic Conclusions trajectory at 0, 6,12 and 24 h Persistent or new congestion within the first 24 hours of cardiogenic shock is associated with worse outcomes.

Higher mortality in SCAI D and E stages



Congestion matters!








Invasive Therapies for Cardiogenic Shock











Early Revascularization in Acute Myocardial Infarction Complicated by Cardiogenic Shock

Authors: Judith S. Hochman, M.D., Lynn A. Sleeper, Sc.D., John G. Webb, M.D., Timothy A. Sanborn, M.D., Harvey D. White, D.Sc., J. David Talley, M.D., Christopher E. Buller, M.D., +16, for the SHOCK Investigators* Author Info & Affiliations





MCS Armamentarium








Intraaortic Balloon Support for Myocardial Infarction with Cardiogenic Shock

Authors: Holger Thiele, M.D., Uwe Zeymer, M.D., Franz-Josef Neumann, M.D., Miroslaw Ferenc, M.D., Hans-Georg Olbrich, M.D., Jörg Hausleiter, M.D., Gert Richardt, M.D., +12, for the IABP-SHOCK II Trial Investigators* Author Info & Affiliations



| Baseline Variable | No. of Patients | IABP | Control | Relative Risk (95% C | :1) | P Value for Interaction |
|---------------------|--------------------|-----------|--------------|----------------------------|------------------|----------------------------|
| | | 30-day mo | ortality (%) | | | |
| Sex | | | | | | 0.61 |
| Female | 187 | 44.4 | 43.2 | | 1.03 (0.74-1.43) | |
| Male | 411 | 37.3 | 40.5 | | 0.92 (0.72-1.18) | |
| Age | | | | | | 0.09 |
| <50 yr | 70 | 19.4 | 44.1 | | 0.44 (0.21-0.95) | |
| 50-75 yr | 334 | 34.6 | 36.5 | | 0.95 (0.71-1.27) | |
| >75 yr | 194 | 53.7 | 50.0 | | 1.07 (0.81-1.41) | |
| Diabetes | | | | | | 0.82 |
| Yes | 195 | 42.9 | 46.7 | | 0.92 (0.67-1.26) | |
| No | 399 | 37.2 | 38.9 | | 0.96 (0.74-1.23) | |
| Hypertension | | | | | | 0.05 |
| Yes | 410 | 42.9 | 40.4 | | 1.06 (0.84-1.34) | |
| No | 183 | 28.9 | 43.0 | | 0.67 (0.45-1.01) | |
| Type of MI | | | | | | 0.76 |
| STEMI/LBBB | 412 | 41.0 | 42.9 | | 0.96 (0.77-1.21) | |
| Non-STEMI | 177 | 37.5 | 38.3 | | 0.98 (0.67-1.43) | |
| STEMI type | | | | | | 0.14 |
| Anterior | 216 | 35.4 | 43.7 | | 0.81 (0.58-1.13) | |
| Nonanterior | 196 | 48.3 | 42.2 | | 1.16 (0.85-1.57) | |
| Previous infarction | | | | | | 0.04 |
| Yes | 131 | 47.9 | 33.3 | + | 1.44 (0.93-2.21) | |
| No | 466 | 37.3 | 43.3 | | 0.86 (0.69-1.07) | |
| Hypothermia | | | | | | 0.31 |
| Yes | 226 | 48.1 | 44.2 | | 1.09 (0.82-1.44) | |
| No | 372 | 35.1 | 39.3 | | 0.89 (0.68-1.16) | |
| Blood pressure | | | | | | 0.76 |
| <80 mm Hg | 161 | 50.7 | 46.4 | | 1.09 (0.79-1.50) | |
| ≥80 mm Hg | 432 | 35.9 | 39.2 | | 0.92 (0.72-1.17) | |
| | | | 0.0 | 0.5 1.0 1.5 2.0 | 2.5 | |
| | | | , | IABP Better Control Better | | |




IABP in HF-CS

CHANGE IN CARDIAC OUTPUT FOLLOWING IABP INSERTION

TABLE 3. Multivariable predictors of hemodynamic response to intra-aortic balloc

| Variable | Odds Ratio |
|---------------------------------------------------------------|------------|
| Heart rate >90 bpm | 1.52 |
| Systemic vascular resistance >1300 dynes/sec/cm ⁻⁵ | 5.04 |
| Pulse pressure >30 mm Hg | 1.07 |
| Right atrium to pulmonary artery diastolic pressure < 0.60 | 1.30 |
| Moderate to severe mitral regurgitation | 2.42 |



Fried JA et al. *J Heart Lung Transplant*. 2018;37(11):1313-1321.

CENTRAL ILLUSTRATION: Early Intra-Aortic Balloon Pump in Heart Failure **Complicated by Cardiogenic Shock** 48 patients Crossover to IABP: 13% randomized for Escalation to other mechanical 101 adult patients with standard care heart failure-cardiogenic shock circulatory support: 4.2% (SoC) (HF-CS) with mean LVEF = 20% Stage B 53 patients randomized for HF-CS intra-aortic balloon pump 28% (IABP) and SoC Escalation to other mechanical Stage C circulatory support: 7.5% HF-CS 57% Survival or Successful Bridge to LVAD or Heart Transplant at 60 Days Probability (%) 50-HR: 0.72 (95% CI: 0.31-1.68) Survival 25-Days After Randomization Number at risk The routine early use of IABP plus standard care, compared to standard care alone, did not result in significantly better survival or successful bridge to LVAD or heart transplantation in patients with HF-CS.

Morici N, et al. JACC. 2025;85(16):1587-1597.

Morici, N et al. JACC. 2025 Apr, 85 (16) 1587–1597.


OCTOBER 11, 2025 • CENTRAL BANK CENTER • LEXINGTON, KY








Microaxial Flow Pump or Standard Care in Infarct-Related Cardiogenic Shock

Authors: Jacob E. Møller, D.M.Sc. , Thomas Engstrøm, D.M.Sc., Lisette O. Jensen, D.M.Sc., Hans Eiskjær, D.M.Sc., Norman Mangner, M.D. , Amin Polzin, M.D., P. Christian Schulze, M.D., +29, for the DanGer Shock Investigators* Author Info & Affiliations















Extracorporeal Life Support in Infarct-Related Cardiogenic Shock

Authors: Holger Thiele, M.D. D., Uwe Zeymer, M.D., Ibrahim Akin, M.D., Michael Behnes, M.D., Tienush Rassaf, M.D., Amir Abbas Mahabadi, M.D., Ralf Lehmann, M.D., +28, for the ECLS-SHOCK Investigators* Author Info & Affiliations








Utility of the Swan-Ganz Catheter



Swan-Ganz Catheter







Original Contribution



Evaluation Study of Congestive Heart Failure and Pulmonary Artery Catheterization EffectivenessThe ESCAPE Trial

The ESCAPE Investigators and ESCAPE Study Coordinators*




Table 4. Primary Outcomes: Mortality and Hospitalizations


| PAC Group | Clinical Assessment Group | End Point Estimate (95% CI)* | χ^2 | <i>P</i> Value |
|--------------|---------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 133 | 135 | Hazard ratio, 1.00 (0.82-1.21) | 0.00 | .99 |
| 141 | 143 | Hazard ratio, 0.99 (0.82-1.21) | 0.00 | .95 |
| 43 | 38 | Odds ratio, 1.26 (0.78-2.03) | 0.86 | .35 |
| 8.7 | 8.3 | Hazard ratio, 1.04 (0.86-1.27) | 0.18 | .67 |
| 0 | 0 | NA | NA | NA |
| 10 | 11 | Odds ratio, 0.97 (0.38-2.22) | 0.04 | .97 |
| | 133 141 43 8.7 | PAC Group Assessment Group 133 135 141 143 43 38 8.7 8.3 0 0 | PAC Group Assessment Group End Point Estimate (95% CI)* 133 135 Hazard ratio, 1.00 (0.82-1.21) 141 143 Hazard ratio, 0.99 (0.82-1.21) 43 38 Odds ratio, 1.26 (0.78-2.03) 8.7 8.3 Hazard ratio, 1.04 (0.86-1.27) 0 0 NA 10 11 Odds ratio, 0.00 | PAC Group Assessment Group End Point Estimate (95% CI)* χ² 133 135 Hazard ratio, 1.00 (0.82-1.21) 0.00 141 143 Hazard ratio, 0.99 (0.82-1.21) 0.00 43 38 Odds ratio, 1.26 (0.78-2.03) 0.86 8.7 8.3 Hazard ratio, 1.04 (0.86-1.27) 0.18 0 0 NA NA 10 11 Odds ratio, 0.04 |

Abbreviations: CI, confidence interval; LVAD, left ventricular assist device; NA, not applicable; PAC, pulmonary artery catheter.




^{*}Values less than 1 favor PAC.

CENTRAL ILLUSTRATION: Frequency of Mortality Among PAC Use Overall and by SCAI Stage



Garan, A.R. et al. J Am Coll Cardiol HF. 2020;8(11):903-13.


CENTRAL ILLUSTRATION: Pulmonary Artery Catheter Use in CICUs and Association With Mortality



Kadosh BS, et al. J Am Coll Cardiol HF. 2023;11(8):e003226.




PACCS trial








24-hour Roadmap to CS management





Sinha, S et al. JACC. 2025 Apr, 85 (16) 1618-1641.

Golden Hour of Acute Cardiogenic Shock





Sinha, S et al. JACC. 2025 Apr, 85 (16) 1618–1641.

Talk 2: Siddarth Pahwa



Cardiogenic shock – A surgeon's perspective

Siddharth V. Pahwa, MD
Associate Professor
Division Chief, Adult Cardiac Surgery
University of Louisville, KY



Disclosures

I do not have any relationships to report with ACCME defined ineligible companies.

I will not be discussing unlabeled/investigational uses of medical devices or pharmaceuticals during this presentation.



Goals of treatment

- Improve abnormal hemodynamics
- Maintain adequate tissue perfusion



Myriad of options available – How do you choose?

- What is the primary problem?
- Which ventricle is significantly involved?



| SHORT-TERM MECHANICAL CIRCULATORY SUPPORT | PERCUTANEOUS OPTIONS | SURGICAL OPTIONS |
|-------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Left-sided | •IABP (femoral or axillary) •Impella 2.5 (LV-Ao) •Impella CP (LV-Ao) •TandemHeart (LA-Ao) | Axillary graft IABP via graft anastomoses Impella 5.0 via axillary graft Impella LD (direct arteriotomy of ascending aorta) LV CentriMag (LV apical position) |
| Right-sided | •Impella RP (RV-PA) •TandemHeart (RA-PA) •ProtekDuo cannula with centrifugal pump (RA-PA) | •RV CentriMag (RV or RA-PA) |
| Biventricular | •Peripheral VA ECMO •Impella CP or 2.5 <i>plus</i> Impella RP •Impella CP or 2.5 <i>plus</i> TandemHeart (RA-PA) or ProtekDuo with centrifugal pump | Central cannulation VA ECMO Biventricular CentriMag |




Left Ventricle

IABP

- Most commonly used modality of temporary MCS
- Decreases diastolic blood pressure, decreases afterload, decreases myocardial oxygen consumption
- Increases coronary perfusion
- IABP Shock II study negative
- True cardiogenic shock No benefit
- Cl 1.8-2 on no inotropes YES!!



Left Ventricle



Impella 5.5

- LV unloading, reduces LVEDP
- Decreases myocardial O2 consumption
- Typically placed via the axillary artery (8 or 10 mm graft)
- Cardiogenic shock involving predominantly the LV or as part of ECPELLA




Left Ventricle

VA ECMO

- Highest level of support
- Inserted percutaneously, or directly via a sternotomy
- Provides 4-6 liters/min flow
- Decompresses RV, increases LV afterload
- In cases with poor LV function (minimal contractility), LV vent is needed
- ECPELLA



Right Ventricle



Protek Duo

- Inserted via the IJV
- Drains blood from the RA and reinfuses into the PA
- Centriimag pump



Case Presentation


- 23 y/o F, peripartum cardiomyopathy, delivered 2 days ago
- Arrested at OSH
- CPR for 8 min, ROSC
- Transferred to us

- On epi 0.1 mics/kg/min, nor epi 0.2 mics/kg/min and vaso at 4 U/min
- Emergent Swan Cl 1.5 l/min/sq m
- TEE RV function moderately reduced, LVEF 5%



- Emergent Impella 5.5 via trans-axillary route
- POD 1 Sinus tach, HR 160-180/min
- TEE RV now severely depressed, Impella 5.5 in good position
- Emergent Protek Duo via RIJ
- Supported on Protek and Impella for 7 days, extubated
- Listed for transplant
- POD 8 Orthotopic Heart Transplant
- POD 16 Discharged home







Questions and Discussion



Talk 2: Jeffrey Spindel



The Future of Critical Care Cardiology

Jeff Spindel, DO



Disclosures

None



Outline

Patient Population

- Diagnoses
- Comorbidities

People

- Training
- Care Models


Systems

- Shock Teams
- Centers






CICU Discharge Diagnosis








Comorbidities





CICU Therapies and Treatment










Changes in patient populations...









...Requires changes in expertise



Training in Critical Care Cardiology Within Critical Care Medicine Fellowship



A Novel Pathway


Connor G. O'Brien, MD, a Christopher F. Barnett, MD, MPH, David M. Dudzinski, MD, JD, Pablo A. Sanchez, MD, Jason N. Katz, MD, MHS, John G. Harold, MD, Erin K. Hennessey, MD, Paul K. Mohabir, MD

- 1. Airway/ventilator management
- 2. Management of mixed shock
- 3. Management of MCS


- 4. Familiarity with renal replacement therapy
- 5. Management of neurologic emergencies
- 6. Postoperative cardiac surgery management















Staffing Models



Primary Physician



Consultations






| Intervention | Mortality | Cost | LOS | Reference |
|---------------------------------|-----------|----------|----------|-----------|
| Closed Unit | | | | 2 |
| HF/CC Staffing | | | | 3,4 |
| 24/7 Staffing | ✓ | | | 5 |
| Mandatory CC Consultation (IMV) | | ✓ | ✓ | 6 |
| Shock Team | ✓ | ✓ | ✓ | 7,8,9,10 |




Shock Teams











Lower mortality


(23 vs 29%), aOR 0.72
 Cost saving¹¹






So, what is the future of CCC?







Systems of Care







The NEW ENGLAND JOURNAL of MEDICIN

JOURNAL OF THE AMERICAN COLLEGE OF CARDIOLOGY © 2018 BY THE AMERICAN COLLEGE OF CARDIOLOGY FOUNDATION

ESTABLISHED IN 1812

OCTOBER 4, 2012

Intraaortic Balloon Support for Myocardial Inf with Cardiogenic Shock

Holger Thiele, M.D., Uwe Zeymer, M.D., Franz-Josef Neumann, M.D., Miroslaw Fe Hans-Georg Olbrich, M.D., Jörg Hausleiter, M.D., Gert Richardt, M.D., Marcus Hennersdorf, M.D. Georg Fuernau, M.D., Steffen Desch, M.D., Ingo Eitel, M.D., Rainer Hambrecht, M.D., Jörg Michael Böhm, M.D., Henning Ebelt, M.D., Steffen Schneider, Ph.D., Gerhard Schuler, M.D., ar for the IABP-SHOCK II Trial Investigators*

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Extracorporeal Life Support in Infarct-Related Cardiogenic Shock

ORIGINAL CONTRIBUTION

Percutaneous Mechanic Epinephrine Versus Norepinephrine

for Cardiogenic Shock After

Supp Pump Acut

> zmar M. xander F rel T. Koo m K. Lagi

tially. However, cardiogenic shock (CS) may occur prior to or following reperfusion. Even those who survive acute intervention may later develop CS and the overall 30-day mortality for patients with CS in association with MI is approximately 40-50%. Unfortunately, this incidence has not changed in the past 20 years since the publication of the landmark SHOCK (SHould we emergently revascularize Occluded Coronaries for cardiogenic shock) trial.²⁻⁵

ras, MD,^c Tristan Morichau-Beauchant, MD,^d Quenot, MD, PhD,^g Antoine Kimmoun, MD, PhD,^a Harjola, MD, PнD, h Ferhat Meziani, MD, PнD, b iarte, РнD, ^j Nicolas Girerd, MD, РнD, ^j



ORIGINAL RESEARCH ARTICLE

JAMA-EXPRESS

Extracorporeal Membrar

The NEW ENGLAND JOURNAL of MEDICINE

Effect

S. Ewer

T. Goslar

and patients aged >75 years. The incidence of CS has increased in recent years, while the reason for increasing incidence is unclear, improved diagnosis and better access to care are both likely contributory. While the in-hospital mortality has improved, the 6- to 12-month mortality in cardiogenic

shock has remained unchanged at ≈50% over the past 2 decades<mark>.³, ⁴</mark>

and Cardiogenic Shock

The TRIUMPH Randomized Controlled Trial

/olovar, MD, Miroslava Seyfrydova, N inhart, MD, PhD, and Jan Belohlavek

ogenic Shock Richard G. Jung, Ph.D.,

th Dobutamine

F. Daniel Ramirez, M.D., David T. Harnett, M.D., Anas Merdad, M.B., B.S., Aws Almufleh, M.B., B.S., Willy Weng, M.D., Omar Abdel-Razek, M.D., Shannon M. Fernando, M.D., Kwadwo Kyeremanteng, M.D., M.H.A., Jordan Bernick, M.Sc., George A. Wells, Ph.D., Vincent Chan, M.D., Michael Froeschl, M.D., C.M., Marino Labinaz, M.D., Michel R. Le May, M.D., Juan J. Russo, M.D., and Benjamin Hibbert, M.D., Ph.D.

Kentucky

OCTOBER 11, 2025 • CENTRAL BANK CENTER • LEXINGTON, KY

Future of Critical Care Cardiology

Patients

- More comorbid
- More non-cardiac illnesses
- More IMV
- More RRT

People

- Expanded CCC interest and training opportunities
- Designated staffing

Systems

- Dedicated Units
- Multidisciplinary Shock Teams
- Interhospital communication and escalation



References

- 1. Miller PE, Chouairi F, Thomas A, Kunitomo Y, Aslam F, Canavan ME, Murphy C, Daggula K, Metkus T, Vallabhajosyula S, Carnicelli A, Katz JN, Desai NR, Ahmad T, Velazquez EJ, Brennan J. Transition From an Open to Closed Staffing Model in the Cardiac Intensive Care Unit Improves Clinical Outcomes. J Am Heart Assoc. 2021 Feb 2;10(3):e018182.
- 2. Sims DB, Kim Y, Kalininskiy A, Yanamandala M, Josephs J, Rivas-Lasarte M, Ahmed N, Assa A, Jahufar F, Kumar S, Sun E, Rahgozar K, Ali SZ, Zhang M, Patel S, Edwards P, Saeed O, Shin JJ, Murthy S, Patel S, Shah A, Jorde UP. Full-Time Cardiac Intensive Care Unit Staffing by Heart Failure Specialists and its Association with Mortality Rates. J Card Fail. 2022 Mar;28(3):394-402.
- 3. Na SJ, Chung CR, Jeon K, Park CM, Suh GY, Ahn JH, Carriere KC, Song YB, Choi JO, Hahn JY, Choi JH, Choi SH, On YK, Gwon HC, Jeon ES, Kim DK, Yang JH. Association Between Presence of a Cardiac Intensivist and Mortality in an Adult Cardiac Care Unit. J Am Coll Cardiol. 2016 Dec 20;68(24):2637-2648.
- 4. Huard P, Kalavrouziotis D, Lipes J, Simon M, Tardif MA, Blackburn S, Langevin S, Sia YT, Mohammadi S. Does the full-time presence of an intensivist lead to better outcomes in the cardiac surgical intensive care unit? J Thorac Cardiovasc Surg. 2020 Apr;159(4):1363-1375.e7.
- 5. Kapoor K, Verceles AC, Netzer G, Chaudhry A, Bolgiano M, Devabhakthuni S, Ludmir J, Pollock JS, Ramani GV, McCurdy MT. A Collaborative Cardiologist-Intensivist Management Model Improves Cardiac Intensive Care Unit Outcomes. J Am Coll Cardiol. 2017 Sep 12;70(11):1422-1423. doi: 10.1016/j.jacc.2017.07.739. Erratum in: J Am Coll Cardiol. 2017 Nov 28;70(21):2737-2738. doi: 10.1016/j.jacc.2017.10.034. PMID: 28882242.
- 6. Tehrani BN, Truesdell AG, Sherwood MW, et al. Standardized team-based care for cardiogenic shock. J Am Coll Cardiol 2019; 73:1659–1669.
- 7. Taleb I, Koliopoulou AG, Tandar A, et al. Shock team approach in refractory cardiogenic shock requiring short-term mechanical circulatory support: a proof of concept. Circulation 2019; 140:98–100.
- 8. Lee F, Hutson JH, Boodhwani M, et al. Multidisciplinary code shock team in cardiogenic shock: A Canadian Centre Experience. CJC Open 2020; 2:249–257.
- 9. Papolos AI, Kenigsberg BB, Berg DD, et al. Management and outcomes of cardiogenic shock in cardiac ICUs with versus without shock teams. J Am Coll Cardiol 2021; 78:1309–1317.
- 10. Taleb I, Giannouchos TV, Kyriakopoulos CP, Clawson A, Davis ES, Sideris K, Tseliou E, Shah KS, Tonna JE, Dranow E, Jones TL, Carter SJ, Fang JC, Stehlik J, Ohsfeldt RL, Selzman CH, Hanff TC, Drakos SG. Cost-Effectiveness of a Shock Team Approach in Refractory Cardiogenic Shock. Circ Heart Fail. 2024 Nov;17(11):e011709. doi:



Jeff Spindel, DO Jeffrey.Spindel@uky.edu



Please evaluate this session using the QR Code



